Summary of product characteristics for a biocidal product

Product name: GMP 34

Product type(s): PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

Authorisation number: 201710-0114

R4BP 3 asset reference number: IS-0021337-0006

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	2
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	3
3. Hazard and precautionary statements	3
4. Authorised use(s)	3
5. General directions for use	8
5.1. Instructions for use	8
5.2. Risk mitigation measures	-
	8
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	8
5.4. Instructions for safe disposal of the product and its packaging	8
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	9
6. Other information	9

1.1. Trade names of the product

Dipal RTU	
Dipal RTU	

1.2. Authorisation holder

Name and address of the	Name DeLaval NV	
authorisation holder	Address	Industriepark-Drongen 10 9031 Gent Belgium
Authorisation number	201710-0114 1-3	
R4BP 3 asset reference number	IS-0021337-0006	

Date of the authorisation

Expiry date of the authorisation

21/08/2019

01/08/2029

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer

Address of the manufacturer

Location of manufacturing sites

DeLaval NV
Industriepark-Drongen 10 9031 Gent Belgium
Industriepark-Drongen 10 9031 Gent Belgium

1.4. Manufacturer(s) of the active substance(s)

Active substance	1319 - Iodine
Name of the manufacturer	Sociedad Quimica y Minera (SQM) S.A.
Address of the manufacturer	Supplier EU: SQM Europe NV - Sint-Pietersvliet 7 bus 8 2000 Antwerpen Belgium
Location of manufacturing sites	Los Militares 4290 Piso 4 8320000 Las Condes (Santiago) Chile

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
lodine		Active Substance	7553-56-2	231-442-4	0,15
Alcohol ethoxylate			68439-46-3		1,25
Sodium iodide			7681-82-5	231-679-3	0,065

2.2. Type of formulation

AL - Any other liquid

3. Hazard and precautionary statements

Hazard statements

Keep out of reach of children.

Precautionary statements

4. Authorised use(s)

4.1 Use description

Use 1 - Teat disinfection, Ready-to-use dip products applied manually

Product type	PT03 - Veterinary hygiene (Disinfectants)
Where relevant, an exact description of the authorised use	Teat disinfectant
Target organism(s) (including development stage)	Scientific name: Common name: Bacteria Development stage:
	Scientific name: Common name: Yeasts Development stage:
	Scientific name: Common name: Viruses Development stage:
Field(s) of use	Indoor
	The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals
Application method(s)	Post-milking teat disinfection of udder teats with ready-to-use products - The product is applied manually after milking on the teats by manual dipping using a dip cup.

Application rate(s) and frequencies	ca. 5 ml per cow; ca. 2.5 ml per sheep or goat - 0% - To be applied after milking, 2 times per day.The products can be used all year during the lactation period.
Category(ies) of users	Professional
Pack sizes and packaging material	Can, Plastic: HDPE , 5L Can, Plastic: HDPE , 10L Can, Plastic: HDPE , 20L Can, Plastic: HDPE , 60L Drum, Plastic: HDPE , 200L IBC, Plastic: HDPE , 1000L

4.1.1 Use-specific instructions for use

The product is a ready-to-use product and is used undiluted. The product must be brought to a temperature above 20°C before use.

Directions for use:

It is recommended to wear gloves during milking and application to protect the skin and ensure hygienic milking condition. For manual application: use a clean dip cup.

Ensure that teats are clean before use. Apply the ready-to-use solution of the product on each teat immediately after milking: DIP: dip every teat in the product. Ensure that the teat is covered to at least three quarters length and replenish the dip cup as necessary;

Allow teats to dry after application during at least 5 min and keep the animals standing during this period.

4.1.2 Use-specific risk mitigation measures

There are no use specific risk mitigation measures. See 5.2 for general instructions.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.2 Use description

-

Use 2 - Teat disinfection, Ready-to-use spray products applied manually

Product type	PT03 - Veterinary hygiene (Disinfectants)	
Where relevant, an exact description of the authorised use	Teat disinfectant	
Target organism(s) (including development stage)		
	Scientific name: Common name: Yeast Development stage:	
	Scientific name: Common name: Viruses Development stage:	
Field(s) of use	Indoor	
()	The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals	
Application method(s)	post-milking teat disinfection of udder teats with ready-to-use products - The product is applied manually after milking on the teats by spraying using a spray container.	
Application rate(s) and frequencies	ca. 5 ml per cow; ca. 2.5 ml per sheep or goat - 0% - To be applied after milking, 2 times per day. The products can be used all year during the lactation period.	
Category(ies) of users	Professional	
Pack sizes and packaging material	Can, Plastic: HDPE , 5L Can, Plastic: HDPE , 10L Can, Plastic: HDPE , 20L Can, Plastic: HDPE , 60L Drum, Plastic: HDPE , 200L IBC, Plastic: HDPE , 1000L	

4.2.1 Use-specific instructions for use

The product is a ready-to-use products and is used undiluted.

The product must be brought to a temperature above 20°C before use.

Directions for use:

For manual application: use a clean applicator.

Ensure that teats are clean before use. Apply the ready-to-use solution of the product on each teat immediately after milking: 1) MANUAL TRIGGER SPRAY: spray every teat 1 sec. with the product. Ensure that the teat is covered to at least three guarters length and replenish the spray container as necessary;

2) AUTOMATIC SPRAYER: spray every teat 1 sec. with the product.

Allow teats to dry after application during at least 5 min and keep the animals standing during this period.

4.2.2 Use-specific risk mitigation measures

Use chemical resistant gloves (glove material to be specified by the authorisation holder within the product information) during postmilking teat disinfection by manual spraying. See 5.2 for general instructions.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

4.3 Use description

Use 3 - Teat disinfection, Ready-to-use dip or spray products applied automatically

Product type

Where relevant, an exact description of the authorised use Scientific name: Target organism(s) (including

development stage)

PT03 - Veterinary hygiene (Disinfectants)	
---	--

Teat disinfectant

Common name: Bacteria Development stage:

Scientific name: Common name: Yeasts Development stage:

	Scientific name: Common name: Viruses Development stage:
Field(s) of use	Indoor
	The product is intended for post-milking teat disinfection. The product is applied directly after milking on the teats of lactating animals
Application method(s)	Post-milking teat disinfection of udder teats with ready-to-use products - The product is applied automatically by milking machines after milking on the teats by integrated dipping or spraying device.
Application rate(s) and frequencies	ca. 5 ml per cow; ca. 2.5 ml per sheep or goat - 0% - To be applied after milking, 3 times per day.The products can be used all year during the lactation period.
Category(ies) of users	Professional
Pack sizes and packaging material	Can, Plastic: HDPE, 5L Can, Plastic: HDPE, 10L Can, Plastic: HDPE, 20L Can, Plastic: HDPE, 60L Drum, Plastic: HDPE, 200L IBC, Plastic: HDPE, 1000L

4.3.1 Use-specific instructions for use

The product is a ready-to-use product and is used undiluted.
The product must be brought to a temperature above 20°C before use

4.3.2 Use-specific risk mitigation measures

There are no use specific risk mitigation measures. See 5.2 for general instructions.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

There are no use specific instructions. See 5.3 for general instructions.

Т

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

There are no use specific instructions. See 5.4 for general instructions.

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

5. General directions for use

5.1. Instructions for use

There are no general instructions for use. See the use specific instructions for use.

5.2. Risk mitigation measures

In case a combination of pre- and post-milking disinfection is necessary, using another product not containing iodine has to be considered for pre-milking disinfection.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

DIRECT/INDIRECT EFFECTS:

According to our experience and to the information provided to us, the product does not have any adverse effects if it is used and handled as specified.

FIRST AID MEASURES:

Eye contact: Rinse thoroughly with plenty of water for at least 15 minutes. If eye irritation persists: Get medical advice/attention.. Skin contact: Wash off immediately with soap and plenty of water removing all contaminated clothes and shoes. Ingestion: Wash out mouth with water. Seek medical advice immediately if symptoms occur and/or large quantities have been ingested. Inhalation: Move to fresh air. EMERGENCY MEASURES TO PROTECT THE ENVIRONMENT: Prevent further leakage or spillage if safe to do so. Methods and materials for containment and clean-up: Dam up. Soak up with inert absorbent material. Prevent product from entering drains. Keep in suitable and closed containers for disposal.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose unused product and the packaging in accordance with local requirements. Used product can be flushed to the municipal sewer or disposed to the manure deposit depending on local requirements. Avoid release to an individual waste water treatment plant.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Storage: Store upright in the tightly closed original container. Protect from direct light, temperatures above 30°C and frost. Shelf-life: 1 year.

6. Other information

The hazard- and precautionary statements shall be in Icelandic on the product label.

The precautionary statement in Icelandic: P102 Geymist þar sem börn ná ekki til.