

Summary of product characteristics for a biocidal product

Product name: Neosorex Bait Blocks

Product type(s): PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

Authorisation number: UK-2012-0360

R4BP 3 asset reference number: UK-0000154-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
4.1.1 Use-specific instructions for use	4
4.1.2 Use-specific risk mitigation measures	4
4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	5
4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	5
4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	5
4.2.1 Use-specific instructions for use	6
4.2.2 Use-specific risk mitigation measures	7
4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	7
4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	7
4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	7
4.3.1 Use-specific instructions for use	8
4.3.2 Use-specific risk mitigation measures	9
4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	9
4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	9
4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	9
4.4.1 Use-specific instructions for use	10
4.4.2 Use-specific risk mitigation measures	10

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	11
4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	11
4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	11
4.5.1 Use-specific instructions for use	12
4.5.2 Use-specific risk mitigation measures	12
4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	12
4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	12
4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	12
4.6.1 Use-specific instructions for use	14
4.6.2 Use-specific risk mitigation measures	14
4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	14
4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging	14
4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage	14
5. General directions for use	14
5.1. Instructions for use	14
5.2. Risk mitigation measures	15
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	16
5.4. Instructions for safe disposal of the product and its packaging	16
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	16
6. Other information	17

Administrative information

1.1. Trade names of the product

Neosorex Bait Blocks

1.2. Authorisation holder

Name and address of the authorisation holder	Name	BASF plc
	Address	Earl Road, Cheadle Hulme, SK8 6QG Cheadle United Kingdom
Authorisation number	UK-2012-0360	
R4BP 3 asset reference number	UK-0000154-0000	
Date of the authorisation	09/08/2012	
Expiry date of the authorisation	10/05/2023	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	BASF Agro B.V., Arnhem (NL) – Freienbach Branch
Address of the manufacturer	Hubostrasse 3 8808 Pfäffikon SZ Zuerich Switzerland
Location of manufacturing sites	BASF plc, St Michael's Industrial, Oldgate WA8 8TJ Widnes United Kingdom

1.4. Manufacturer(s) of the active substance(s)

Active substance	26 - Difenacoum
Name of the manufacturer	BASF Agro B.V. Arnhem (NL) - Freienbach Branch
Address of the manufacturer	Hubostrasse 3 8808 Pfäffikon SZ Zuerich Switzerland
Location of manufacturing sites	Vertellus Specialities UK Limited, Lower Road, Halebank, WA8 8NS Widnes United Kingdom

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Difenacoum	3-(3-biphenyl-4-yl-1,2,3,4-tetrahydro-1-naphthyl)-4-hydroxycoumarin	Active Substance	56073-07-5	259-978-4	0.005

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements	<p>May damage the unborn child.</p> <p>May cause damage to organs (blood) through prolonged or repeated exposure .</p>
Precautionary statements	<p>Obtain special instructions before use.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Wear protective gloves.</p> <p>IF exposed or concerned: Get medical advice.</p> <p>Get medical advice if you feel unwell.</p> <p>Store locked up.</p>

Dispose of container to in accordance with national regulations.

4. Authorised use(s)

4.1 Use description

Use 1 - Professional users with demonstrated competence (equivalent to trained professionals) – indoor

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Mus musculus-House mouse-Adult and juveniles.

Rattus norvegicus-Brown rat-Adults and juveniles

Rattus rattus-Black rat or roof rat-Adults and juveniles

Field(s) of use

Indoor

Indoor

Application method(s)

Bait application -

- Ready-to-use bait to be used in tamper-resistant bait stations

- Covered and protected baiting points

Application rate(s) and frequencies

See below - 100 -

Rats:

- up to 200g of bait per baiting point placed 10 metres apart.

Mice:

- up to 30g of bait per baiting point placed 1-2 metres apart.

Category(ies) of users

Trained professional

Pack sizes and packaging material

- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg
- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for mice containing either loose blocks or paper/laminate sachets (30g) – 3kg to 25kg
- Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg
- Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg
- Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.1.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.

UK only: Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting"

4.1.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Do not use this product in pulsed baiting treatments.

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient

- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

4.2 Use description

Use 2 - Professional users with demonstrated competence (equivalent to trained professionals) – outdoor around buildings

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Rattus norvegicus-Brown rat-Adults and juveniles
Rattus rattus-Black rat / roof rat-Adults and juveniles
Mus musculus-House mouse-Adults and juveniles

Field(s) of use

Outdoor
Outdoor around buildings

Application method(s)

Bait application -
- Ready-to-use bait to be used in tamper-resistant bait stations .
- Covered and protected baiting points
- Direct application methods to burrows

Application rate(s) and frequencies

See below - 100 -

Rats:

- up to 200g g of bait per baiting point placed 10 metres apart.

Mice:

- up to 30g g of bait per baiting point placed 1-2 metres apart.

Burrow baiting: up to 200g of bait per burrow.

Category(ies) of users

Trained professional

Pack sizes and packaging material

- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg
- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for mice containing either loose blocks or paper/laminate sachets (30g) – 3kg to 25kg
- Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg
- Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg
- Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.2.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas non-labile to flooding.
 - Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
 - For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
 - Remove the remaining product at the end of treatment period (except when directly applied into burrows).
 - Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent).
- UK only: Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on

Permanent Baiting"

- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population. [When available] Follow any additional instructions provided by the relevant code of best practice.

4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the applicable code of good practice.

- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Do not use this product in pulsed baiting treatments.

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient

- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

4.3 Use description

**Use 3 - Professional users with demonstrated competence (equivalent to trained professionals) –
Outdoor open areas & waste dumps**

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Rattus norvegicus-Brown rat-Adults and juveniles Rattus rattus-Roof rat-Adults and juveniles
Field(s) of use	Outdoor Outdoor open areas Outdoor waste dumps
Application method(s)	Bait application - - Ready-to-use bait to be used in tamper-resistant bait stations . - Covered and protected baiting points - Direct application methods to burrows
Application rate(s) and frequencies	See below - 100 - - Up to 200g of bait per baiting point placed 10 metres apart. Burrow baiting: up to 200g of bait per burrow.
Category(ies) of users	Trained professional
Pack sizes and packaging material	<ul style="list-style-type: none"> • Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg • Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg • Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg • Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas non-labile to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Remove the remaining product at the end of treatment period (except when directly applied into burrows).
- Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows

to reduce the risks of bait being rejected and spilled. Follow any additional instructions provided by the CRRU UK Code of Best Practice (or equivalent).
 UK only: Sites under a permanent baiting regime should be inspected regularly in accordance with product label directions. The period between visits should be determined by the technician in charge but will not be longer than every four weeks when permanent baiting is conducted outdoors. For permanent baiting follow any additional instructions provided by the CRRU Guidance on Permanent Baiting”
 - Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population. [When available] Follow any additional instructions provided by the relevant code of best practice.

4.3.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign in accordance with the applicable code of good practice.
 - To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
 - Do not use this product in pulsed baiting treatments.
 - Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient
 - The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

4.4 Use description

Use 4 - Professional users with demonstrated competence (equivalent to trained professionals) – sewers

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Rattus norvegicus-Brown rat-Adults and juveniles

Field(s) of use	Other Sewers
Application method(s)	Bait application - - Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water. - Covered and protected baiting points
Application rate(s) and frequencies	See below - 100 - Up to 200g per manhole.
Category(ies) of users	Trained professional
Pack sizes and packaging material	<ul style="list-style-type: none"> • Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg • Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg • Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg • Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.4.1 Use-specific instructions for use

- Baits must be applied in a way so that they do not come into contact with water and are not washed away.

4.4.2 Use-specific risk mitigation measures

- Do not use this product in pulsed baiting treatments.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See "Directions for use"

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

4.5 Use description

Use 5 - Professional users – indoor – NOT RELEVANT IN UK

Product type

PT14 - Rodenticides (Pest control)

Where relevant, an exact description of the authorised use

Not relevant for rodenticides

Target organism(s) (including development stage)

Rattus norvegicus-Brown rat-Adults and juveniles
Mus musculus-House mouse-Adults and juveniles
Rattus rattus-Roof rat-Adults and juveniles

Field(s) of use

Indoor
Indoor

Application method(s)

Bait application -
Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

See below - 100 -
Rats:
- up to 200g of bait per baiting point placed 10 metres apart.
Mice:
- up to 30g of bait per baiting point placed 1-2 metres apart.

Category(ies) of users

Professional

Pack sizes and packaging material

- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg
- Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for mice containing either loose blocks or paper/laminate sachets (30g) – 3kg to 25kg
- Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg
- Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg
- Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.5.1 Use-specific instructions for use

- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.5.2 Use-specific risk mitigation measures

See "Directions for use"

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

4.6 Use description

Use 6 - Professional users – outdoor around buildings – NOT RELEVANT IN UK

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Mus musculus-House mouse-Adults and juveniles Rattus norvegicus-Brown rat-Adults and juveniles Rattus rattus-Roof rat-Adults and juveniles
Field(s) of use	Outdoor Outdoor around buildings
Application method(s)	Bait application - Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequencies	See below - 100 - Rats: - up to 200g of bait per baiting point placed 10 metres apart. Mice: - up to 30g of bait per baiting point placed 1-2 metres apart.
Category(ies) of users	Professional
Pack sizes and packaging material	<ul style="list-style-type: none">• Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for rats containing either loose blocks or paper/laminate sachets (200g) – 3kg to 25kg• Overpacked pre-formed and sealed cardboard/plastic bait box or tray with peelable lid for mice containing either loose blocks or paper/laminate sachets (30g) – 3kg to 25kg• Polypropylene/HDPE container or similar container such as lidded bucket, pot or bottle with resealable lid containing either loose blocks or paper/laminate sachets (containing either 4g or 23g blocks) – 3kg to 25kg• Paper-laminated/poly-weave or similar resealable bag containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) – 3kg to 25kg• Lined re-closable cardboard carton containing either loose blocks or paper/laminate sachets (containing 4g or 23g blocks) - 3kg to 10kg

4.6.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas non-labile to flooding.
- For mice: The baiting stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- For rats: The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.
- [When available] Follow any additional instructions provided by the relevant code of best practice.

4.6.2 Use-specific risk mitigation measures

- Do not apply this product directly in the burrows.

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See "Directions for use"

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See "Directions for use"

5. General directions for use

5.1. Instructions for use

Professional users with demonstrated competence (equivalent to trained professionals)

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by CRRU UK Code of Best Practice (or equivalent). Sites where burrow baiting is used should be visited more frequently than those where secure bait boxes are employed.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- For non-emptiable sachets: Do not open the sachets containing the bait].

Professional users – NOT RELEVANT IN UK

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 7.3 for the information to be shown on the label).
- [If national policy or legislation require it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

Professional users with demonstrated competence (equivalent to trained professionals)

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
 - Do not use in areas where resistance to the active substance can be suspected.
 - Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment (unless permanent baiting).
 - Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
 - Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
 - Dispose dead rodents in accordance with local requirements
- (Not relevant in the UK) -The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only)
- UK only: In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).
- UK only: To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.
- UK only: Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.
- UK only: When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

Professional users – NOT RELEVANT IN UK

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign.
- To reduce risk of secondary poisoning, search for and remove dead rodents at frequent intervals during treatment (e.g. at least twice a week).
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment.
- Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").
- Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed) the user should seek advice from the product supplier or call a pest control service.
- Do not wash the bait stations with water between applications.
- Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
 - Antidote: Vitamin K1 administered by medical/veterinary personnel only.
 - In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.
- UK only: UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. Contact a veterinary surgeon in case of ingestion by a pet.
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison center [insert national phone number]".
 - Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose uneaten bait and the packaging in accordance with local requirements.
 - Professional users with demonstrated competence (equivalent to trained professionals) only: Use of gloves is recommended.
- UK only: For information on disposal in the UK contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months.

6. Other information

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

This product contains a bittering agent and a dye.